

A RELAXED EVALUATION SUBGROUP

TOSHIHIRO YAMAGUCHI

ABSTRACT. Let $f : X \rightarrow Y$ be a pointed map between connected CW-complexes. As a generalization of the evaluation subgroup $G_*(Y, X; f)$, we will define the *relaxed evaluation subgroup* $\mathcal{G}_*(Y, X; f)$ in the homotopy group $\pi_*(Y)$ of Y , which is identified with $\text{Im}\pi_*(\tilde{e}v)$ for the evaluation map $\tilde{e}v : \text{map}(X, Y; f) \times X \rightarrow Y$ given by $\tilde{e}v(h, x) = h(x)$. Especially we see by using Sullivan model in rational homotopy theory for the rationalized map $f_{\mathbb{Q}}$ that $\mathcal{G}_*(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = \pi_*(Y) \otimes \mathbb{Q}$ if the map f induces an injection of rational homotopy groups. Also we compare it with more relaxed subgroups by several rationalized examples.

1. INTRODUCTION

Let $f : X \rightarrow Y$ and $g : B \rightarrow Y$ be pointed maps of connected CW complexes. Recall the definition of pairing with axes f and g [O], which is given by the existence of a map $F_{g,f} : B \times X \rightarrow Y$ in the homotopy commutative diagram:

$$(1) \quad \begin{array}{ccc} B \times X & \xleftarrow{i_X} & X \\ i_B \uparrow & \searrow F_{g,f} & \downarrow f \\ B & \xrightarrow{g} & Y, \end{array}$$

where $i_B(b) = (b, *)$ and $i_X(x) = (*, x)$. In particular, when $B = Y = X$ and $f = g = id$, X is an H-space of the multiplication $F_{id,id}$. In this paper, we consider whether or not there exist a section $s : B \rightarrow B \times X$ and a map $F : B \times X \rightarrow Y$ such that the diagram

$$(2) \quad \begin{array}{ccc} B \times X & \xleftarrow{i_X} & X \\ s \uparrow & \searrow F & \downarrow f \\ B & \xrightarrow{g} & Y. \end{array}$$

homotopically commutes. Here a section means a map $s : B \rightarrow B \times X$ which satisfies $p_B \circ s \simeq id_B$ for the projection $p_B : B \times X \rightarrow B$ with $p_B(b, x) = b$.

The n th Gottlieb group $G_n(X)$ of a space X is the subgroup of the n th homotopy group $\pi_n(X)$ of X consisting of homotopy classes of maps $a : S^n \rightarrow X$ such that the wedge $(a|id_X) : S^n \vee X \rightarrow X$ extends to a map $F_a : S^n \times X \rightarrow X$ [G]. The n th evaluation subgroup $G_n(Y, X; f)$ of a map $f : X \rightarrow Y$ is the subgroup of $\pi_n(Y)$ represented by maps $a : S^n \rightarrow Y$ such that $(a|f) : S^n \vee X \rightarrow Y$ extends to a map

MSC: 55P62, 55Q70, 55R15

Keywords: Gottlieb group, evaluation subgroup, relaxed evaluation subgroup, tncz subgroup, sectional subgroup, rational homotopy, Sullivan model, derivation

$F_{a,f} : S^n \times X \rightarrow Y$ inducing the homotopy commutative diagram

$$(3) \quad \begin{array}{ccc} S^n \times X & \xleftarrow{i_X} & X \\ i_{S^n} \uparrow & \searrow F_{a,f} & \downarrow f \\ S^n & \xrightarrow{a} & Y \end{array}$$

[WK], which is the case of $B = S^n$ in (1). The map $F_{a,f}$ in (3) is the adjoint of a map $F'_{a,f}$ in the homotopy commutative diagram

$$(3)' \quad \begin{array}{ccc} & & \text{map}_f(X, Y) \\ & \nearrow F'_{a,f} & \downarrow ev \\ S^n & \xrightarrow{a} & Y \end{array}$$

where $F'_{a,f}(b)(x) := F_{a,f}(b, x)$ for $b \in S^n$ and $x \in X$. Here $\text{map}_f(X, Y)$ is the connected component of f in the mapping space $\text{map}(X, Y)$ from X to Y with the compact open topology and ev is the evaluation map given by $ev(h) = h(*)$. Then

$$G_n(Y, X; f) = \text{Im} (\pi_n(ev) : \pi_n(\text{map}_f(X, Y)) \rightarrow \pi_n(Y))$$

in $\pi_n(Y)$ and therefore it is called an ‘evaluation’ subgroup of a map. Notice that (3) is a special case of the homotopy commutative diagram

$$(4) \quad \begin{array}{ccc} S^n \times X & \xleftarrow{i_X} & X \\ \eta \uparrow & \searrow F & \downarrow f \\ S^n & \xrightarrow{a} & Y, \end{array}$$

in which $\eta : X \xrightarrow{i_X} S^n \times X \xrightarrow{p_{S^n}} S^n$ is a trivial fibration with a section $s : S^n \rightarrow S^n \times X$. Here we can put $s(b) = (b, \tau(b))$ for a map $\tau : S^n \rightarrow X$.

Definition A. For a map $f : X \rightarrow Y$, the n th *relaxed evaluation subgroup* of f is given as

$$\begin{aligned} \mathcal{G}_n(Y, X; f) &:= \{a \in \pi_n(Y) \mid \text{there are a section } s : S^n \rightarrow S^n \times X \\ &\text{and a map } F : S^n \times X \rightarrow Y \text{ such that } F \circ s \simeq a, f \simeq F \circ i_X\}. \end{aligned}$$

The map F in (4) is the adjoint of a map $\tilde{F}'$ in the homotopy commutative diagram

$$(4)' \quad \begin{array}{ccc} & & \text{map}_f(X, Y) \times X \\ & \nearrow \tilde{F}' & \downarrow \tilde{ev} \\ S^n & \xrightarrow{a} & Y \end{array}$$

where $\tilde{F}'(b) := (F'(b), \tau(b))$ with $F'(b)(x) = F(b, x)$ and $s(b) = (b, \tau(b))$ for $b \in S^n$ and $x \in X$. Here $\tilde{ev} : \text{map}(X, Y; f) \times X \rightarrow Y$ is the evaluation map given by $\tilde{ev}(h, x) = h(x)$. Thus

Claim. $\mathcal{G}_n(Y, X; f) = \text{Im}(\pi_n(\tilde{e}v) : \pi_n(\text{map}_f(X, Y) \times X) \rightarrow \pi_n(Y))$.

In this paper, we will estimate $\text{Im}\pi_n(\tilde{e}v)$ in several cases according to Definition A. We note that it is a naturally generalized object of an ordinary evaluation subgroup. Indeed, for a subcomplex X' of X , we can put

$$\mathcal{G}_n(Y, X; f)(X') := \{a \in \pi_n(Y) \mid \text{there are a section } s : S^n \rightarrow S^n \times X' \text{ and a map } F : S^n \times X \rightarrow Y \text{ such that } F \circ s \simeq a, f \simeq F \circ i_X\}.$$

Then we have

$$\mathcal{G}_n(Y, X; f) \supset \mathcal{G}_n(Y, X; f)(X') \supset \mathcal{G}_n(Y, X; f)(*) = G_n(Y, X; f)$$

and $\mathcal{G}_n(Y, X; f)(X') = \text{Im}(\pi_n(\tilde{e}v) : \pi_n(\text{map}_f(X, Y) \times X') \rightarrow \pi_n(Y))$. In the following, we see several properties of a relaxed evaluation subgroup.

Lemma 1.1. *For a subspace X of a space Y and a map $g : B \rightarrow Y$ with $\text{Im } g \subset X$, there are a section $s : B \rightarrow B \times X$ and a map $F : B \times X \rightarrow X$ such that the diagram*

$$\begin{array}{ccc} B \times X & \xleftarrow{i_X} & X \\ \uparrow s & \searrow F & \downarrow \cap \\ B & \xrightarrow{g} & Y \end{array}$$

commutes.

Indeed, put $s(b) = (b, g(b))$ and $F((b, x)) = x$ for $(b, x) \in B \times X$.

In particular, in the case of $B = S^n$ and $X = Y^m$, the m -skelton of Y , we have by cellular approximation theorem

Proposition 1.2. *For $n \leq m$, $\mathcal{G}_n(Y, Y^m; i_{Y^m}) = \pi_n(Y)$ for $i_{Y^m} : Y^m \subset Y$. In particular, $\mathcal{G}_*(Y, Y; id_Y) = \pi_*(Y)$.*

Of course $\mathcal{G}_n(Y, *, *) = G_n(Y, *, *) = \pi_n(Y)$ for the constant map $* : * \rightarrow Y$ and we know $G_n(Y) = G_n(Y, Y; id_Y) \subset G_n(Y, X; f)$ for any map $f : X \rightarrow Y$. In contrast, $\mathcal{G}_n(Y, X; f) \subset \mathcal{G}_n(Y, Y; id_Y)$ from Proposition 1.2. Note that $\mathcal{G}_*(Y, X; f)$ may be zero for a map $f : X \rightarrow Y$ even if $\pi_*(Y) \neq 0$ (see Example 2.6 and examples in Section 3).

Also as an evaluation subgroup satisfies it, a relaxed evaluation subgroup satisfies

$$\pi_n(g)(\mathcal{G}_n(Y, X; f)) \subset \mathcal{G}_n(Y', X; g \circ f)$$

for a pointed map $g : Y \rightarrow Y'$. Thus there is a map $\pi_n(f) : \pi_n(X) = \mathcal{G}_n(X, X; id_X) \rightarrow \mathcal{G}_n(Y, X; f)$. Therefore the relaxed evaluation subgroups $\mathcal{G}_n(X, Z; j)$ and $\mathcal{G}_n(Y, X; f)$ are embedded into parts of the homotopy exact sequence of a fibration $\xi : Z \xrightarrow{j} X \xrightarrow{f} Y$, respectively.

Corollary 1.3. *For a fibration $\xi : Z \xrightarrow{j} X \xrightarrow{f} Y$ and any n , there are the sequences*

$$\pi_{n+1}(Y) \xrightarrow{\partial_{n+1}} \pi_n(Z) \xrightarrow{\pi_n(j)} \mathcal{G}_n(X, Z; j) \xrightarrow{\pi_n(f)} \pi_n(Y)$$

and

$$\pi_n(Z) \xrightarrow{\pi_n(j)} \pi_n(X) \xrightarrow{\pi_n(f)} \mathcal{G}_n(Y, X; f) \xrightarrow{\partial_n} \pi_{n-1}(Z)$$

which are both exact. Moreover, for the connecting map $\tilde{\partial} : \Omega Y \rightarrow Z$ of ξ ,

$$\pi_{n+1}(X) \xrightarrow{\pi_{n+1}(f)} \pi_{n+1}(Y) \xrightarrow{\partial_{n+1}} \mathcal{G}_n(Z, \Omega Y; \tilde{\partial}) \xrightarrow{\pi_n(j)} \pi_n(X)$$

is exact.

Note that, for a pointed map $g : X' \rightarrow X$, there is an inclusion $G_n(Y, X; f) \subset G_n(Y, X'; f \circ g)$. But it does not hold for relaxed evaluation subgroups in general.

Lemma 1.4. *For a map $g : B \rightarrow Y$ and a map $f : X \rightarrow Y$ such that $f_* : [B, X] \rightarrow [B, Y]$ is surjective, there are a section $s : B \rightarrow B \times X$ and a map $F : B \times X \rightarrow X$ such that the diagram*

$$\begin{array}{ccc} B \times X & \xleftarrow{i_X} & X \\ \uparrow s & \searrow F & \downarrow f \\ B & \xrightarrow{g} & Y \end{array}$$

homotopically commutes.

Indeed, there is a lift $\tilde{g} : B \rightarrow X$ such that $f \circ \tilde{g} \simeq g$ from the assumption. Then put $s(b) = (b, \tilde{g}(b))$ and $F((b, x)) = f(x)$ for $(b, x) \in B \times X$.

In particular, in the case of $B = S^n$, we have

Proposition 1.5. *If a map $f : X \rightarrow Y$ induces a surjection $\pi_n(f) : \pi_n(X) \rightarrow \pi_n(Y)$, then $\mathcal{G}_n(Y, X; f) = \pi_n(Y)$.*

For example, $\mathcal{G}_*(Y, X; f) = \pi_*(Y)$ if a map f has a section. In contrast, $G_*(Y, X; f) = G_*(Y)$ if f has a section.

Suppose that $f_{\mathbb{Q}} : X_{\mathbb{Q}} \rightarrow Y_{\mathbb{Q}}$ is a rationalized map with X and Y simply connected CW complexes of finite type [HMR]. We consider the relaxed evaluation subgroup of a map and more relaxed subgroups from a point of view of rational homotopy. By using Sullivan's minimal model arguments, we show

Theorem 1.6. *If a map $f : X \rightarrow Y$ induces an injection $\pi_*(f) \otimes \mathbb{Q} : \pi_*(X) \otimes \mathbb{Q} \rightarrow \pi_*(Y) \otimes \mathbb{Q}$ on rational homotopy groups, then $\mathcal{G}_*(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = \pi_*(Y) \otimes \mathbb{Q}$.*

From Proposition 1.5 and Theorem 1.6, we have

Corollary 1.7. *If the homotopy fiber of a map $f : X \rightarrow Y$ has the rational homotopy type of the Eilenberg-MacLane space $K(\mathbb{Q}, n)$ for some n , then $\mathcal{G}_*(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = \pi_*(Y) \otimes \mathbb{Q}$.*

In Section 2, we prove Theorem 1.6 after preparing of the notion of derivations of model. In Section 3, we will define more relaxed subgroups of the homotopy group of Y , the *tncz subgroup* $\mathcal{T}_*(Y, X; f)$ and the *sectional subgroup* $\mathcal{S}_*(Y, X; f)$, for a map $f : X \rightarrow Y$. They are defined by relaxing the trivial fibration η in (4). We compare $\mathcal{G}_*(Y, X; f)$ with them by several rationalized examples.

2. SULLIVAN MODELS

We use the Sullivan minimal model $M(Y)$ of a simply connected CW complex Y of finite type. It is a free $\mathbb{Q}$ -commutative differential graded algebra (DGA) $(\Lambda W, d_Y)$ with a $\mathbb{Q}$ -graded vector space $W = \bigoplus_{i \geq 2} W^i$ where $\dim W^i < \infty$ and a decomposable differential. Here $\Lambda^+ W$ is the ideal of ΛW generated by elements of positive degree. Denote the degree of a homogeneous element x of a graded algebra as $|x|$. Then $xy = (-1)^{|x||y|}yx$ and $d(xy) = d(x)y + (-1)^{|x|}xd(y)$. Recall $M(Y)$ determines the rational homotopy type of Y . Especially there is an isomorphism

$W^i \cong \text{Hom}(\pi_i(Y), \mathbb{Q})$. We denote the dual element of $a \in \pi_i(Y) \otimes \mathbb{Q}$ as a^* . Put $M(Y) = (\Lambda W, d_Y)$. Then the model of a map $f : X \rightarrow Y$ is given by a KS-extension

$$(\Lambda W, d_Y) \xrightarrow{i} (\Lambda W \otimes \Lambda V, D) \xrightarrow{q} (\Lambda V, \overline{D})$$

with $D|_{\Lambda W} = d_Y$ and the minimal model $(\Lambda V, \overline{D})$ of the homotopy fiber of f or

$$(H^*(Y; \mathbb{Q}), 0) \xrightarrow{i} (\Lambda W \otimes \Lambda V, D) \xrightarrow{q} (\Lambda V, \overline{D})$$

when Y is formal (for example, when Y is a sphere) [FHT]. In general, D is not decomposable and it is decomposable if and only if $\pi_*(f) \otimes \mathbb{Q} : \pi_*(X) \otimes \mathbb{Q} \rightarrow \pi_*(Y) \otimes \mathbb{Q}$ is a surjection. See [FHT] for a general introduction and the standard notations.

Let A be a DGA $A = (A^*, d_A)$ with $A^* = \oplus_{i \geq 0} A^i$, $A^0 = \mathbb{Q}$, $A^1 = 0$ and the augmentation $\epsilon : A \rightarrow \mathbb{Q}$. Define $\text{Der}_i A$ the vector space of self-derivations of A decreasing the degree by $i > 0$, where $\theta(xy) = \theta(x)y + (-1)^{|x|}x\theta(y)$ for $\theta \in \text{Der}_i A$. We denote $\oplus_{i > 0} \text{Der}_i A$ by $\text{Der} A$. The boundary operator $\delta : \text{Der}_* A \rightarrow \text{Der}_{*-1} A$ is defined by $\delta(\sigma) = d_A \circ \sigma - (-1)^{|\sigma|} \sigma \circ d_A$. For a DGA-map $\phi : A \rightarrow B$, define a ϕ -derivation of degree n to be a linear map $\theta : A^* \rightarrow B^{*-n}$ with $\theta(xy) = \theta(x)\phi(y) + (-1)^{|x|} \phi(x)\theta(y)$ and $\text{Der}(A, B; \phi)$ the vector space of ϕ -derivations. The boundary operator $\delta_\phi : \text{Der}_*(A, B; \phi) \rightarrow \text{Der}_{*-1}(A, B; \phi)$ is defined by $\delta_\phi(\sigma) = d_B \circ \sigma - (-1)^{|\sigma|} \sigma \circ d_A$. Note $\text{Der}_*(A, A; id_A) = \text{Der}_*(A)$. For $\phi : A \rightarrow B$, the composition with $\epsilon' : B \rightarrow \mathbb{Q}$ induces a chain map $\epsilon'_* : \text{Der}_n(A, B; \phi) \rightarrow \text{Der}_n(A, \mathbb{Q}; \epsilon)$. For a minimal model $A = (\Lambda Z, d_A)$, define $G_n(A, B; \phi) := \text{Im}(H(\epsilon'_*) : H_n(\text{Der}(A, B; \phi)) \rightarrow \text{Hom}_n(Z, \mathbb{Q}))$. Especially $G_*(A, A; id_A) = G_*(A)$. Note that $z^* \in \text{Hom}(Z, \mathbb{Q})$ (z^* is the dual of the basis element z) is in $G_n(A, B; \phi)$ if and only if z^* extends to a derivation $\theta \in \text{Der}(A, B; \phi)$ with $\delta_\phi(\theta) = 0$.

Theorem 2.1. ([FH],[LS]) $G_n(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) \cong G_n(M(Y), M(X); M(f)) = G_n((\Lambda W, d_Y), (\Lambda W \otimes \Lambda V, D))$.

Thus $G_n(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}})$ is completely determined by the derivations only. For a DGA-map $\phi : (\Lambda V, d) \rightarrow (\Lambda Z, d')$, the symbol $(v, h) \in \text{Der}(\Lambda V, \Lambda Z; \phi)$ means the ϕ -derivation sending an element $v \in V$ to $h \in \Lambda Z$ and the other to zero. Especially $(v, 1) = v^*$.

Example 2.2. Consider the fibration $S^3 \rightarrow X \xrightarrow{f} Y = S^2 \times S^2$ whose KS-extension is given by

$$(\Lambda(w_1, w_2, w_3, w_4), d_Y) \rightarrow (\Lambda(w_1, w_2, w_3, w_4, v), D) \rightarrow (\Lambda(v), 0),$$

where $|w_1| = |w_2| = 2$, $|w_3| = |w_4| = |v| = 3$, $d_Y w_1 = d_Y w_2 = 0$, $d_Y w_3 = w_1^2$, $d_Y w_4 = w_2^2$ and $Dv = w_1 w_2$. Since D is decomposable, $\pi_*(f) \otimes \mathbb{Q} : \pi_*(X) \otimes \mathbb{Q} \rightarrow \pi_*(Y) \otimes \mathbb{Q}$ is surjective. So we have $\mathcal{G}_2(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = \pi_2(Y) \otimes \mathbb{Q}$ from Proposition 1.5. On the other hand, from Theorem 2.1, $G_2(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = 0$ since

$$\delta_f((w_i, 1)) = 2(w_{i+2}, w_i) \notin \delta_f(\text{Der}(\Lambda W, \Lambda W \otimes \Lambda^+ v))$$

for $i = 1, 2$ and $W = \mathbb{Q}(w_1, w_2, w_3, w_4)$. Note that $f_{\mathbb{Q}}$ has no section ([T]).

Proof of Theorem 1.6. Fix an element $a \in \pi_n(Y) \otimes \mathbb{Q}$. From the assumption, there is a DGA-projection $p_{W,V} : (\Lambda W, d_Y) \rightarrow (\Lambda V, \overline{d}_Y)$ as the model of f . A model of

the (non homotopy commutative) diagram

$$\begin{array}{ccc} E & \xleftarrow{i} & X_{\mathbb{Q}} \\ p \downarrow & & \downarrow f_{\mathbb{Q}} \\ S_{\mathbb{Q}}^n & \xrightarrow{a} & Y_{\mathbb{Q}} \end{array}$$

is given by

$$\begin{array}{ccc} (\Lambda x/x^2 \otimes \Lambda V, D') & \xrightarrow{q} & (\Lambda V, \overline{d}_Y) \\ \cup \uparrow & & \uparrow p_{W,V} \\ (\Lambda x/x^2, 0) & \xleftarrow{M(a)} & (\Lambda W, d_Y) \end{array}$$

where $|x| = n$, $i_{V,W} : V \subset W$ and $p_{W,V}(\Lambda W) = \overline{\Lambda W} = \Lambda V$ with $p_{W,V} \circ i_{V,W} = id_V$. Here $\Lambda x/x^2 = \Lambda x$ if n is odd and $\Lambda x/x^2 = \mathbb{Q}[x]/(x^2)$ if n is even.

We will construct a rationally trivial fibration of the form $\eta : X_{\mathbb{Q}} \rightarrow E \rightarrow S_{\mathbb{Q}}^n$ together with a suitable map $F : E \rightarrow Y_{\mathbb{Q}}$, in model terms. Define a graded $\mathbb{Q}$ -algebra map $F' : \Lambda W \rightarrow \Lambda x/x^2 \otimes \Lambda V$ by

$$F'(w) = \overline{w} + (-1)^{n|w|}\sigma(w)x$$

where $\sigma \in Der_n(\Lambda W, \Lambda V; p_{W,V})$ with $(-1)^{n|u|}\sigma(u)x = M(a)(u)$ for $u \in W$. Also define the differential D' by $D'(x) = 0$ and

$$D'|_{\Lambda V} = \overline{d}_Y - \delta_{\overline{d}_Y}(\overline{\sigma}) \cdot x,$$

where $\overline{\sigma} \in Der_n(\Lambda V)$ is uniquely given by $\overline{\sigma}(\overline{w}) = \sigma(w)$ for $w \in \Lambda W$ and $(\theta \cdot x)(z) := (-1)^{n|z|}\theta(z)x$ for a derivation θ . Then $D' \circ D' = 0$ from $\delta_{\overline{d}_Y} \circ \delta_{\overline{d}_Y} = 0$ and F' is a DGA-map by

$$\begin{aligned} F'd_Y(w) &= \overline{d_Y(w)} + (-1)^{n(|w|+1)}\sigma(d_Y w)x \\ &= \overline{d_Y(w)} - (-1)^{n|w|}\delta_{\overline{d}_Y}(\overline{\sigma})(\overline{w})x + (-1)^{n|w|}\overline{d}_Y\sigma(w)x \\ &= D'(\overline{w} + (-1)^{n|w|}\sigma(w)x) = D'F'(w) \end{aligned}$$

for $w \in \Lambda W$. Thus we have the KS-model of η

$$(\Lambda x/x^2, 0) \xrightarrow{i} (\Lambda x/x^2 \otimes \Lambda V, D') \xrightarrow{q} (\Lambda V, \overline{d}_Y)$$

and a map

$$F' : (\Lambda W, d_Y) \rightarrow (\Lambda x/x^2 \otimes \Lambda V, D').$$

Since $\delta_{\overline{d}_Y}(\overline{\sigma}) \in Der(\Lambda V, \Lambda^+ V)$, the fibration η has a section s ([T]). Moreover the definition of D' indicates the rational triviality of η :

$$(\Lambda x/x^2 \otimes \Lambda V, D') \cong (\Lambda x/x^2, 0) \otimes (\Lambda V, \overline{d}_Y)$$

over $(\Lambda x/x^2, 0)$ since then the homotopy class of the classifying map $S^n \rightarrow Baut_1 X_{\mathbb{Q}}$, $[\delta_{\overline{d}_Y}(\overline{\sigma})]$, is zero in $\pi_n(Baut_1 X) \otimes \mathbb{Q} = H_{n-1}(Der M(X))$ [S]. We

can choose the model of s by $M(s)(x) = x$ and $M(s)(z) = 0$ for $z \in \Lambda V$, then $M(s) \circ F' = M(a)$. Thus there is a DGA-commutative diagram

$$\begin{array}{ccc} (\Lambda x/x^2 \otimes \Lambda V, D') & \xrightarrow{q} & (\Lambda V, \bar{d}_Y) \\ M(s) \downarrow & \swarrow F' & \uparrow p_{W,V} \\ (\Lambda x/x^2, 0) & \xleftarrow{M(a)} & (\Lambda W, d_Y). \end{array}$$

It is the rational model of (4). $\square$

Proof of Corollary 1.7. Put the model of homotopy fiber $(\Lambda v, 0)$. When Dv is decomposable, we have from Proposition 1.5. Also when Dv is not decomposable, there is a surjection $M(Y) = (\Lambda W, d_Y) \rightarrow M(X) = (\Lambda V, d_X)$ with $\dim V = \dim W - 1$. Then we have from Theorem 1.6. $\square$

Remark 1. In the proof of above, the fibration $\eta : X_{\mathbb{Q}} \xrightarrow{i} E \xrightarrow{p} S_{\mathbb{Q}}^n$ is trivial, that is, there is a homotopy commutative diagram

$$\begin{array}{ccc} E & \xleftarrow{i} & X_{\mathbb{Q}} \\ p \downarrow & \swarrow g & \downarrow i_{X_{\mathbb{Q}}} \\ S_{\mathbb{Q}}^n & \xleftarrow{p_{S_{\mathbb{Q}}^n}} & S_{\mathbb{Q}}^n \times X_{\mathbb{Q}} \end{array}$$

where g is a homotopy equivalence. The model $M(g)$ is given by $id - \bar{\sigma} \cdot x$, which is a quasi-isomorphism. But the map g does not induce the homotopy commutative diagram with a section s of p

$$\begin{array}{ccc} E & \xleftarrow{i} & X_{\mathbb{Q}} \\ s \uparrow & \swarrow g & \downarrow i_{X_{\mathbb{Q}}} \\ S_{\mathbb{Q}}^n & \xrightarrow{i_{S_{\mathbb{Q}}^n}} & S_{\mathbb{Q}}^n \times X_{\mathbb{Q}} \end{array}$$

in general. Therefore, even if $\pi_*(f) \otimes \mathbb{Q}$ is surjective or injective, we can not induce $G_*(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = \pi_*(Y) \otimes \mathbb{Q}$ in general. For example, see Example 2.2 or Example 2.5, respectively.

In the followings, we consider some examples, whose models are given by $M(s)(x) = x$ and $M(s)(z) = 0$ for $z \in M(X)$ as in the proof of Theorem 1.6. The index of an element means the degree.

Example 2.3. For $n > 0$, $\mathcal{G}_n(S_{\mathbb{Q}}^n, S_{\mathbb{Q}}^n; id_{S_{\mathbb{Q}}^n}) = \pi_n(S^n) \otimes \mathbb{Q}$ is given by the following commutative diagrams with $|x| = n$.

$$\begin{array}{ccc} (n : \text{odd}) & (\Lambda x/x^2 \otimes \Lambda w_n, 0) & \xrightarrow{q} (\Lambda w_n, 0) \\ M(s) \downarrow & \swarrow F & \uparrow = \\ (\Lambda x/x^2, 0) & \xleftarrow{M(a)} & (\Lambda w_n, 0) \end{array}$$

where $M(s)(x) = x$, $M(s)(w_n) = 0$ and $F(w_n) = w_n + cx$ for $M(a)(w_n) = cx$ with $c \in \mathbb{Q}$.

$$\begin{array}{ccc}
 (n : \text{even}) & (\Lambda x/x^2 \otimes \Lambda(w_n, w_{2n-1}), D') & \xrightarrow{q} (\Lambda(w_n, w_{2n-1}), d_Y) \\
 & \downarrow M(s) & \uparrow = \\
 & (\Lambda x/x^2, 0) & \xleftarrow{M(a)} (\Lambda(w_n, w_{2n-1}), d_Y)
 \end{array}$$

$\swarrow F$

where $d_Y w_n = 0$, $d_Y w_{2n-1} = w_n^2$, $D'w_n = 0$, $D'w_{2n-1} = w_n^2 + 2cw_n x$, $F(w_n) = w_n + cx$ and $F(w_{2n-1}) = w_{2n-1}$.

Example 2.4. For the Hopf fibration $S^3 \rightarrow S^7 \xrightarrow{f} S^4$, we know $G_4(S_{\mathbb{Q}}^4, S_{\mathbb{Q}}^7; f_{\mathbb{Q}}) = \pi_4(S^4) \otimes \mathbb{Q}$ [LS2]. Indeed, the KS-extension is given by

$$(\Lambda(w_4, w_7), d_Y) \rightarrow (\Lambda(w_4, w_7, v_3), D) \rightarrow (\Lambda(v_3), 0)$$

with $d_Y w_4 = 0$, $d_Y w_7 = w_4^2$ and $Dv_3 = w_4$. Then, from Theorem 2.1, $G_4(S_{\mathbb{Q}}^4, S_{\mathbb{Q}}^7; f_{\mathbb{Q}}) = G_4((\Lambda(w_4, w_7), d_Y), (\Lambda(w_4, w_7, v_3), D)) = \mathbb{Q}(w_4^*)$ since $\delta_f((w_4, 1) - 2(w_7, v_3)) = 0$. Thus we have

$$G_4(S_{\mathbb{Q}}^4, S_{\mathbb{Q}}^7; f_{\mathbb{Q}}) = \mathcal{G}_4(S_{\mathbb{Q}}^4, S_{\mathbb{Q}}^7; f_{\mathbb{Q}}) = \pi_4(S^4) \otimes \mathbb{Q} \cong \mathbb{Q}.$$

Also for the product fibration $S^3 \rightarrow X = S^7 \times S^4 \xrightarrow{f \times id} S^4 \times S^4 = Y$ with the trivial fibration $* \rightarrow S^4 \rightarrow S^4$, we have $G_4(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; (f \times id)_{\mathbb{Q}}) \cong \mathbb{Q}$ and $\mathcal{G}_4(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; (f \times id)_{\mathbb{Q}}) = \pi_4(Y) \otimes \mathbb{Q} \cong \mathbb{Q} \oplus \mathbb{Q}$.

Example 2.5. Suppose that the model of a map $f : X \rightarrow Y$ is given by the projection

$$p_{W,V} : M(Y) = (\Lambda(w_3, w_5, w_7, w_9), d_Y) \rightarrow (\Lambda(w_3, w_7, w_9), \bar{d}_Y) = M(X)$$

where $p_{W,V}(w_5) = \bar{w}_5 = 0$, $d_Y w_3 = d_Y w_5 = 0$, $d_Y w_7 = w_3 w_5$, $d_Y w_9 = w_3 w_7$ and $\bar{d}_Y w_3 = \bar{d}_Y w_7 = 0$ and $\bar{d}_Y w_9 = w_3 w_7$. Then we have $G_3(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = G_7(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = 0$ by the direct calculations of derivations. But we have $\mathcal{G}_*(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = \pi_*(Y) \otimes \mathbb{Q}$ from Theorem 1.6.

Example 2.6. When $f : S^3 \times S^3 \rightarrow S^6$ is the map collapsing $S^3 \vee S^3$, then we show

$$\mathcal{G}_6(S^6, S^3 \times S^3; f) = 0.$$

By degree arguments, any fibration $\eta : S^3 \times S^3 \xrightarrow{i} E \xrightarrow{p} S^6$ is rationally trivial, especially $D = 0$ in the KS-extension

$$(\mathbb{Q}[w_6]/(w_6^2), 0) \rightarrow (\mathbb{Q}[w_6]/(w_6^2) \otimes \Lambda(u_3, v_3), D) \rightarrow (\Lambda(u_3, v_3), 0),$$

where $H^*(S^6; \mathbb{Q}) = \mathbb{Q}[w_6]/(w_6^2)$ and $M(S^3 \times S^3) = (\Lambda(u_3, v_3), 0)$. In particular $E_{\mathbb{Q}} \simeq (S^6 \times S^3 \times S^3)_{\mathbb{Q}}$.

For $a \neq 0 \in \pi_6(S^6)$, suppose that there exists a map $F : E \rightarrow S^6$ in

$$\begin{array}{ccc}
 E & \xleftarrow{i} & S^3 \times S^3 \\
 \downarrow s & \searrow F & \downarrow f \\
 S^6 & \xrightarrow{a} & S^6,
 \end{array}$$

Then

$$M(F_{f,g})(w_6) = cx + u_3v_3$$

for some non-zero $c \in \mathbb{Q}$ associated to a . Then $w_6^2 = 0$ in $H^{12}(S^6; \mathbb{Q})$ but

$$[M(F_{f,g})(w_6)]^2 = [2cxu_3v_3] \neq 0$$

in $H^{12}(E; \mathbb{Q}) = H^{12}(S^6 \times S^3 \times S^3; \mathbb{Q})$. It is a contradiction.

3. THE OTHER SUBGROUPS

Futher we will relax Definition A. For pointed maps $f : X \rightarrow Y$ and $g : B \rightarrow Y$, consider whether or not there exists the homotopy commutative diagram

$$(5) \quad \begin{array}{ccc} E & \xleftarrow{i} & X \\ \downarrow s & \searrow F & \downarrow f \\ B & \xrightarrow{g} & Y \end{array}$$

in which $\eta : X \xrightarrow{i} E \xrightarrow{p} B$ is a fibration with the section s .

Recall that a fibration $Z \xrightarrow{i} X \xrightarrow{p} Y$ is said to be *tncz* (totally non-cohomologous to zero) if $H^*(X) \cong H^*(Z) \otimes H^*(Y)$ as $H^*(Y)$ -modules by $p^* : H^*(Y) \rightarrow H^*(X)$.

Definition B. For a map $f : X \rightarrow Y$, the n th *tncz subgroup* of f is given as

$$\mathcal{T}_n(Y, X; f) := \{a \in \pi_n(Y) \mid \text{there are a tncz fibration } \eta : X \xrightarrow{i} E \xrightarrow{p} S^n \text{ with} \\ \text{a section } s \text{ and a map } F : E \rightarrow Y \text{ such that } F \circ s \simeq a, f \simeq F \circ i\}.$$

Definition C. For a map $f : X \rightarrow Y$, the n th *sectional subgroup* of f is given as

$$\mathcal{S}_n(Y, X; f) := \{a \in \pi_n(Y) \mid \text{there are a fibration } \eta : X \xrightarrow{i} E \xrightarrow{p} S^n \text{ with} \\ \text{a section } s \text{ and a map } F : E \rightarrow Y \text{ such that } F \circ s \simeq a, f \simeq F \circ i\}.$$

Remark 2. Note that $\mathcal{S}_n(Y, X; f)$ is a group. For $a, b \in \mathcal{S}_n(Y, X; f)$, there is a homotopy commutative diagram

$$\begin{array}{ccccc} E_{a+b} & \xrightarrow{h} & E_a \cup_X E_b & \xleftarrow{i} & X \\ \downarrow & \text{pull-back} & \downarrow & \searrow G & \downarrow f \\ S^n & \xrightarrow{q} & S^n \vee S^n & \xrightarrow{(a|b)} & Y, \end{array}$$

where q is the pinching comultiplication and the dotted arrow G is given by the universality of push-out from $F_{a,f} : E_a \rightarrow Y$ and $F_{b,f} : E_b \rightarrow Y$. A section $s : S^n \rightarrow E_{a+b}$ is given by $s(x) := (x, s'q(x))$ for a section $s' : S^n \vee S^n \rightarrow E_a \cup_X E_b$ with $G \circ s' \simeq (a|b)$. Also $i_{a+b} : X \rightarrow E_{a+b}$ is given by the universality of pull-back

from $i : X \rightarrow E_a \cup_X E_b$ and $*$: $X \rightarrow S^n$. Thus we have a homotopy commutative diagram

$$\begin{array}{ccc} E_{a+b} & \xleftarrow{i_{a+b}} & X \\ s \uparrow & \searrow G \circ h & \downarrow f \\ S^n & \xrightarrow{a+b} & Y. \end{array}$$

That is $a + b := (a|b) \circ q \in \mathcal{S}_n(Y, X; f)$. When $\eta_a : X \rightarrow E_a \rightarrow S^n$ and $\eta_b : X \rightarrow E_b \rightarrow S^n$ are trivial (tncz), the fibration $X \rightarrow E_{a+b} \rightarrow S^n$ is trivial (tncz). Thus $\mathcal{G}_n(Y, X; f)$ ($\mathcal{T}_n(Y, X; f)$) is a group too.

We have the sequence of inclusions of groups:

$$G_n(f) \subset \mathcal{G}_n(f) \subset \mathcal{T}_n(f) \subset \mathcal{S}_n(f) \subset \pi_n(Y)$$

for a map $f : X \rightarrow Y$.

We consider them under some conditions on $X_{\mathbb{Q}}$ for a map $f : X \rightarrow Y$. For the KS-extension of the fibration $\eta : X \rightarrow E \rightarrow S^n$ in (5), we see $D' - d_X = 0$ if $\pi_{\geq n}(X) \otimes \mathbb{Q} = 0$. Thus

Proposition 3.1. *If $\pi_{\geq n}(X) \otimes \mathbb{Q} = 0$, $\mathcal{S}_n(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = G_n(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}})$.*

For example, $G_6(f_{\mathbb{Q}}) = \mathcal{G}_6(f_{\mathbb{Q}}) = \mathcal{T}_6(f_{\mathbb{Q}}) = \mathcal{S}_6(f_{\mathbb{Q}}) = 0$ for the map f in Example 2.6. We know that a fibration $X \rightarrow E \rightarrow B$ is homotopically trivial if the classifying map $B \rightarrow \text{Baut}_1 X$ is homotopic to the constant map. Therefore we have

Proposition 3.2. *If $\pi_n(\text{Baut}_1 X) \otimes \mathbb{Q} = 0$, $\mathcal{G}_n(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}}) = \mathcal{S}_n(Y_{\mathbb{Q}}, X_{\mathbb{Q}}; f_{\mathbb{Q}})$.*

In Example 3.4 and Example 3.5 below, $\pi_n(\text{Baut}_1 X) \otimes \mathbb{Q} = H_{n-1}(\text{Der} M(X)) \neq 0$. But it is known that any fibration with fiber $\mathbb{C}P^m$, the m dimensional complex projective space, is rationally tncz. In general, we note

Lemma 3.3. *For $n > 1$, any fibration with fiber X over S^n is rationally tncz if and only if the map $\rho : H_{n-1}(\text{Der} M(X)) \rightarrow \text{Der}_{n-1} H^*(X; \mathbb{Q})$ with $\rho([\sigma])([w]) = [\sigma(w)]$ is zero. Here $\text{Der}_{n-1} H^*(X; \mathbb{Q})$ means the derivations of the graded algebra $H^*(X; \mathbb{Q})$ decreasing the degree by $n-1$ ([FOT, 9.7.2]).*

Proof. The KS-extension of a fibration $X \rightarrow E \rightarrow S^n$ is given by the differential $Dv = dv + \sigma(v)x$ for some $[\sigma] \in H_{n-1}(\text{Der} M(X))$ with the differential d of $M(X)$ and $v \in M(X)$. Then an element $[w]$ of $H^*(X; \mathbb{Q})$ is extend to an element $[w + w'x]$ of $H^*(E; \mathbb{Q})$ if and only if $dw' = \sigma(w)$. $\square$

Example 3.4. For the associated fibration $S^2 \rightarrow \mathbb{C}P^3 \xrightarrow{f} S^4$ of the Hopf fibration $S^3 \rightarrow S^7 \xrightarrow{f} S^4$, the KS-model is given by

$$(\Lambda(w_4, w_7), d_Y) \rightarrow (\Lambda(w_4, w_7, v_2, v_3), D) \rightarrow (\Lambda(v_2, v_3), d)$$

where $d_Y w_4 = 0$, $d_Y w_7 = w_4^2$, $Dv_2 = dv_2 = 0$, $dv_3 = v_2^2$ and $Dv_3 = v_2^2 - w_4$. Also $M(\mathbb{C}P^3) \cong (\Lambda(v_2, w_7), d_X)$ with $d_X v_2 = 0$, $d_X w_7 = v_2^4$ and then $M(f)(w_4) = v_2^2$, $M(f)(w_7) = w_7$. Then we have

$$\mathcal{T}_4(S_{\mathbb{Q}}^4, \mathbb{C}P_{\mathbb{Q}}^3; f_{\mathbb{Q}}) = \pi_4(S^4) \otimes \mathbb{Q} = \mathbb{Q}.$$

Indeed, for $a \in \pi_4(S^4) \otimes \mathbb{Q}$ with $M(a)(w_4) = cx$ ($c \in \mathbb{Q}$) and $M(a)(w_7) = 0$, put

$$D'v_2 = 0, \quad D'w_7 = v_2^4 + 2cv_2^2x$$

$$\text{and } F(w_4) = v_2^2 + cx, \quad F(w_7) = w_7$$

in

$$\begin{array}{ccc} (\Lambda x/x^2 \otimes \Lambda(v_2, w_7), D') & \longrightarrow & (\Lambda(v_2, w_7), d_X) \\ \downarrow & \nwarrow F & \uparrow M(f) \\ (\Lambda x/x^2, 0) & \xleftarrow{M(a)} & (\Lambda(w_4, w_7), d_Y). \end{array}$$

Thus $a \in \mathcal{T}_4(S_{\mathbb{Q}}^4, \mathbb{C}P_{\mathbb{Q}}^3; f_{\mathbb{Q}})$. On the other hand, $\mathcal{G}_4(S_{\mathbb{Q}}^4, \mathbb{C}P_{\mathbb{Q}}^3; f_{\mathbb{Q}}) = 0$ since $(\Lambda x/x^2 \otimes \Lambda(v_2, w_7), D')$ can not be isomorphic to $(\Lambda x/x^2, 0) \otimes (\Lambda(v_2, w_7), d_X)$ over $(\Lambda x/x^2, 0)$ for any D' .

Example 3.5. For the map $f : \mathbb{C}P^2 \rightarrow S^4$ collapsing the 2-cell, $M(f) : M(S^4) = (\Lambda(w_4, w_7), d_Y) \rightarrow (\Lambda(v_2, v_5), d_X) = M(\mathbb{C}P^2)$ is given by $M(f)(w_4) = v_2^2$ and $M(f)(w_7) = v_2v_5$. Then we have

$$\mathcal{T}_4(S_{\mathbb{Q}}^4, \mathbb{C}P_{\mathbb{Q}}^2; f_{\mathbb{Q}}) = \pi_4(S^4) \otimes \mathbb{Q} = \mathbb{Q}.$$

Indeed, for $a \in \pi_4(S^4) \otimes \mathbb{Q}$ with $M(a)(w_4) = cx$ ($c \in \mathbb{Q}$) and $M(a)(w_7) = 0$, put

$$D'(v_2) = 0, \quad D'(v_5) = v_2^3 + 2cv_2x$$

$$\text{and } F(w_4) = v_2^2 + cx, \quad F(w_7) = v_2v_5$$

in

$$\begin{array}{ccc} (\Lambda x/x^2 \otimes \Lambda(v_2, v_5), D') & \longrightarrow & (\Lambda(v_2, v_5), d_X) \\ \downarrow & \nwarrow F & \uparrow M(f) \\ (\Lambda x/x^2, 0) & \xleftarrow{M(a)} & (\Lambda(w_4, w_7), d_Y). \end{array}$$

Thus $a \in \mathcal{T}_4(S_{\mathbb{Q}}^4, \mathbb{C}P_{\mathbb{Q}}^2; f_{\mathbb{Q}})$. On the other hand, we have $\mathcal{G}_4(S_{\mathbb{Q}}^4, \mathbb{C}P_{\mathbb{Q}}^2; f_{\mathbb{Q}}) = 0$ since $(\Lambda x/x^2 \otimes \Lambda(v_2, v_5), D')$ can not be isomorphic to $(\Lambda x/x^2, 0) \otimes (\Lambda(v_2, v_5), d_X)$ over $(\Lambda x/x^2, 0)$ for any D' .

Example 3.6. Put $\xi : \Omega X \xrightarrow{i} LX \xrightarrow{p} X$ the fibration of free loops, in which ΩX is the loop space and $LX = \text{map}(S^1, X)$ is the free loop space of a simply connected space X . It has the section $s : X \rightarrow LX$ with $s(z)$ the constant loop at a point z in X . Consider the case that $X = S^2$. Then $\mathcal{S}_2(LS^2, \Omega S^2; i) \ni s$ since we can choose $F = id_{LS^2}$ as

$$\begin{array}{ccc} LS^2 & \xleftarrow{i} & \Omega S^2 \\ \uparrow s & \searrow = & \downarrow i \\ S^2 & \xrightarrow{s} & LS^2. \end{array}$$

Thus we have $\mathcal{S}_2(LS^2, \Omega S^2; i) \neq 0$. Especially, we see $\mathcal{S}_2(LS_{\mathbb{Q}}^2, \Omega S_{\mathbb{Q}}^2; i_{\mathbb{Q}}) \neq 0$ since s is the torsion free generator of $\pi_2(LS^2)$.

But $\mathcal{T}_2(LS_{\mathbb{Q}}^2, \Omega S_{\mathbb{Q}}^2; i_{\mathbb{Q}}) = 0$. Indeed, the KS-model of ξ is given by

$$(\Lambda(x, y), d_Y) \rightarrow (\Lambda(x, y, \bar{x}, \bar{y}), D) \rightarrow (\Lambda(\bar{x}, \bar{y}), 0)$$

where $M(S^2) = (\Lambda(x, y), d_Y)$ with $@|x| = 2$, $|y| = 3$, $d_Y x = 0$, $d_Y y = x^2$, $|\bar{x}| = 1$, $|\bar{y}| = 2$, $D(\bar{x}) = 0$, and $D(\bar{y}) = 2x\bar{x}$ [VS]. For the KS-model of a fibration $\eta : \Omega S^2 \xrightarrow{i} E \xrightarrow{p} S^2$ with a section is given as

$$(\Lambda(x, y), d_Y) \rightarrow (\Lambda(x, y, \bar{x}, \bar{y}), D') \rightarrow (\Lambda(\bar{x}, \bar{y}), 0)$$

where $D'(\bar{x}) = 0$ and $D'(\bar{y}) = cx\bar{x}$ for some $c \in \mathbb{Q}$ by the degree arguments. If $c = 0$, there does not exist a map $F : (\Lambda(x, y, \bar{x}, \bar{y}), D) \rightarrow (\Lambda(x, y, \bar{x}, \bar{y}), D') = (\Lambda(x, y, \bar{x}, \bar{y}), d_Y)$ that we want. If $c \neq 0$, it is not rationally tncz since $H^*(\Omega S^2; \mathbb{Q}) = \Lambda(\bar{x}, \bar{y})$ and

$$H^*(E; \mathbb{Q}) \cong \mathbb{Q}[x, \{u_i\}_{i>0}] \otimes \Lambda(\bar{x}) / (x^2, x\bar{x}, \{xu_i\}, \{\bar{x}u_i\}, \{u_i u_j\}),$$

where $u_i = [\bar{x}\bar{y}^i]$.

Example 3.7. Put $\xi : S^1 \xrightarrow{j} K \xrightarrow{f} S^1$ the fiber bundle with total space a Klein bottle K . Then $G_1(K, S^1, j) = \mathcal{G}_1(K, S^1, j) = \mathcal{T}_1(K, S^1, j) = \mathbb{Z}$ and $\mathcal{S}_1(K, S^1, j) = \pi_1(K)$.

REFERENCES

- [FH] Y.Félix and S.Halperin, *Rational LS category and its applications*, Trans. A.M.S. **273** (1982) 1-38
- [FHT] Y.Félix, S.Halperin and J.-C.Thomas, *Rational homotopy theory*, Springer G.T.M. **205** [2001].
- [FOT] Y.Félix, J.Oprea and D.Tanré, *Algebraic models in geometry*, Oxford **17** [2008]
- [G] D.H.Gottlieb, *Evaluation subgroups of homotopy groups*, Amer. J. Math. **91** (1969) 729-756
- [HMR] P.Hilton, G.Mislin and J.Roitberg, *Localization of nilpotent groups and spaces*, North-Holland Math. Studies **15** (1975)
- [LS] G.Lupton and S.B.Smith, *Rationalized evaluation subgroups of a map. I. Sullivan models, derivations and G-sequences*, J. Pure Appl. Algebra **209** (1) (2007) 159-171
- [LS2] G.Lupton and S.B.Smith, *The evaluation subgroup of a fibre inclusion*, Topology and its Applications **154** (2007) 1107-1118
- [O] N.Oda, *Pairings and copairings in the category of topological spaces*, Publ. Res. Inst. Math. Sci. Kyoto Univ. **28** (1992) 83-97
- [S] D.Sullivan, *Infinitesimal computations in topology*, Publ. IHES **47** (1978) 269-331
- [T] J.-C. Thomas, *Rational homotopy of Serre fibrations*, Ann. Inst. Fourier, Grenoble **31** (1981) 71-90
- [VS] M.Vigué-Poirrier and D.Sullivan, *The homology theory of the closed geodesic problem*, ibid. **11** (1976) 633-644
- [WK] M.H.Woo and J.R.Kim, *Certain subgroups of homotopy groups*, J.Korean Math. Soc. **21** (1984) 109-120

FACULTY OF EDUCATION, KOCHI UNIVERSITY, 2-5-1, KOCHI, 780-8520, JAPAN
E-mail address: tyamag@kochi-u.ac.jp